Author Index

Alton, K.B.
—, Petruzzi, R.F. and Patrick, J.E.
High-performance liquid chromatographic assay for dilevalol in human plasma and urine using a PRP-1 column and fluorimetric detection 363
Ashraf, M., see Singh, A.K. 245
Awni, W.M.
— and Maloney, J.A.
Optimized high-performance liquid chromatographic method for the analysis of cyclosporine and three of its metabolites in blood and urine 233
Bailey, E.
—, Peal, J.A. and Philbert, M.
Determination of 1,3-dinitrobenzene and its metabolites in rat blood by capillary gas chromatography with electron-capture detection 187
Brandin, S., see Polacek, J. 424
Breuer, J., see Haas, R.H. 47
Brundrett, R.B., see Chen, T.-L. 303
Cache, Ph., see Capon, C. 35
Capon, C.
—, Cache, Ph., Leroy, Y., Strecker, G., Montreuil, J. and Fournet, B.
Isolation of the major O-glycosidically linked oligosaccharides obtained by alkaline borohydride degradation of human meconium glycoproteins 35
Carlin, A.S.
—, Prasad, V.K., Sager, A.O., Simmons, J.E. and Skelly, J.P.
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase high-performance liquid chromatography 162
Castoldi, D.
—, Monzani, V. and Tofanetti, O.
Simultaneous determination of nimesulide and hydroxynimesulide in human plasma and urine by high-performance liquid chromatography 413
Chan, K.
Comparison of gas chromatographic and high-performance liquid chromatographic assays for the determination of debrisoquine and its 4-hydroxy metabolite in human fluids 311
Chen, T.-L.
—, Grochow, L.B., Hurowitz, L.A. and Brundrett, R.B.
Determination of busulfan in human plasma by gas chromatography with electron-capture detection 303
Collins-Nakai, R.L., see Pereira, C.M. 208
Csipo, I., see Nagy, E. 214
Czejka, M.J.
— and Georgopoulos, A.
Mitoxantron determination using high-performance liquid chromatography: improved sensitivity by loop-column injection for dual-dose pharmacokinetic studies 429
Damm, P., see Lehr, K.-H. 153
Daniel, R.M., see Patchett, M.L. 269
Dareer, S.M. El, see Rose, L.M. 419
Degrell, I., see Nagy, E. 214
De Maack, F., see Guichardant, M. 25
Diquet, B., see Joulin, Y. 402
Drozd, J., see Koupil, P. 99
Dumont, M., see Lemaire, S. 77
Eckhardt-Wallasch, H., see Wallasch, Th. M. 175
Edlund, P.O.
Determination of coenzyme Q_{10}, α-tocopherol and cholesterol in biological samples by coupled-column liquid chromatography with coulometric and ultraviolet detection 87
El Dareer, S.M., see Rose, L.M. 419
Eller, T.D., see Vrbanac, J.J. 1
Fournet, B., see Capon, C. 35
Fox, K.E., see Sullivan, W.R. 396
Fr[illegible], F., see Graf, P. 353
Fujimoto, T., see Tawa, R. 143
Furuta, T., see Hirota, N. 237
Georgopoulos, A., see Czejka, M.J. 429
Gercken, G., see Schatowitz, B. 257
Gmur, D.J.
—, Meier, P. and Yee, G.C.
High-performance liquid chromatographic column-switching method for two cyclosporine metabolites in blood 343
Gomaa, K.S., see Marzouki, Z.M.H. 11
Goodwin, K.R., see Schmerr, M.J.F. 379
Gordon, M.J., see Huang, X. 385
Goto, J.
—, Sano, Y., Tsuchiya, K. and Nambara, T.
Studies on steroids. CCXXXIII. Separation and characterization of bile acid disul-

phates in human urine by high-performance liquid chromatography 59
Graf, P.
—, Frueh, F. and Schmid, K.
Determination of the angiotensin converting enzyme inhibitor benazeprilat in plasma and urine by an enzymic method 353
Grochow, L.B., see Chen, T.-L. 303
Guichardant, M.
—, Lagarde, M., Lesieur, M. and De Maack, F.
Thermospray-mass spectrometric analysis of underivatized monohydroxy fatty acids: application to stimulated platelets 25
Gunnarsson, P.O., see Polacek, J. 424
Gupta, R.N.
— and Lew, M.
Determination of isoniazid in plasma by liquid chromatography 441
Haas, R.H.
—, Breuer, J. and Hammen, M.
High-performance liquid chromatographic measurement of selected blood citric acid cycle intermediates 47
Hammen, M., see Haas, R.H. 47
Henning, K., see Wallasch, Th. M. 175
Hill, D.L., see Rose, L.M. 419
—, see Waud, W.R. 220
Hirose, S., see Tawa, R. 143
Hirota, N.
—, Furuta, T. and Kasuya, Y.
Determination of cortisol in human plasma by capillary gas chromatography–mass spectrometry using [$^2H_5$]cortisol as an internal standard 237
Horai, Y., see Kubota, K. 67
Huang, X.
—, Gordon, M.J. and Zare, R.N.
Quantitation of Li^+ in serum by capillary zone electrophoresis with an on-column conductivity detector 385
Hung, C.T.
—, Lim, J.K.C., Zoest, A.R. and Lam, F.C.
Optimization of high-performance liquid chromatographic analysis for isoxazolyl penicillins using factorial design 331
Hurowitz, L.A., see Chen, T.-L. 303
Idle, J.R., see Schmid, B.J. 107
Iri, H., see Matsumoto, K. 323
Ishizaki, T., see Kubota, K. 67
Jamali, F., see Mehvar, R. 135
Jehl, F., see Koechlin, C. 197
Johno, I., see Suzuki, K. 435
Joulin, Y.
—, Moretti, J.L. and Diquet, B.
Highly sensitive method for determination of isopropyl-*p*-iodoamphetamine by gas chromatography with electron-capture detection 402
Jung, S.S.
— and Wittliff, J.L.
Influence of column length and pore size on high-performance ion-exchange chromatography of estrogen and progestin receptors 293
Kasuya, Y., see Hirota, N. 237
Kikuchi, H., see Matsumoto, K. 323
Kitazawa, S., see Suzuki, K. 435
Knapp, D.R., see Vrbanac, J.J. 1
Koechlin, C.
—, Jehl, F. and Monteil, H.
Determination of PI_A and PII_A, the two main components of pristinamycin, by high-performance liquid chromatography in plasma 197
Kojima, K.
—, Uezono, Y., Takahashi, T. and Nakanishi, Y.
High-performance liquid chromatographic method for the determination of a cyclandelate metabolite, mandelic acid, in human plasma 203
Koshide, K., see Tawa, R. 143
Koupil, P.
—, Novák, J. and Drozd, J.
Comparison of the absolute calibration method with the method of standard addition for the determination of halothane in blood by gas chromatographic headspace analysis 99
Kováčová, D., see Svobodová, X. 391
Kubota, K.
—, Horai, Y., Kushida, K. and Ishizaki, T.
Determination of benzoic acid and hippuric acid in human plasma and urine by high-performance liquid chromatography 67
Kuhn, W., see Wallasch, Th.M. 175
Kusala, S., see Svobodová, X. 391
Kushida, K., see Kubota, K. 67
Lagarde, M., see Guichardant, M. 25
Lam, F.C., see Hung, C.T. 331
Lange, U., see Wallasch, Th. M. 175
Lehr, K.-H.
— and Damm, P.
Quantification of the enantiomers of ofloxacin in biological fluids by high-performance liquid chromatography 153
Lemaire, S.
—, Dumont, M. and Nolet, S.

Sensitive method of detection, quantitation and purification of peptides using pre-column derivatization with phenyl isothiocyanate 77
Leroy, Y., see Capon, C. 35
Lesieur, M., see Guichardant, M. 25
Lew, M., see Gupta, R.N. 441
Lim, J.K.C., see Hung, C.T. 331
Luhmann, I., see Szathmary, S.Cs. 193
Maack, F. De, see Guichardant, M. 25
Maaten, M.J. Van Der, see Schmerr, M.J.F. 379
Maloney, J.A., see Awni, W.M. 233
Marzouki, Z.M.H.
—, Taha, A.M. and Gomaa, K.S.
Fatty acid profiles of sebaceous triglycerides by capillary gas chromatography with mass-selective detection 11
Matsumoto, K.
—, Kikuchi, H., Iri, H., Takahasi, H. and Umino, M.
Automated determination of drugs in serum by column-switching high-performance liquid chromatography. II. Separation of theophylline and its metabolites 323
Mehvar, R.
—, Jamali, F. and Pasutto, F.M.
Rapid and sensitive high-performance liquid chromatographic assay of tiaprofenic acid enantiomers in human plasma and urine 135
Meier, P., see Gmur, D.J. 343
Miller, C., see Slegowski, M.B. 227
Monk, C.R., see Patchett, M.L. 269
Monteil, H., see Koechlin, C. 197
Montreuil, J., see Capon, C. 35
Monzani, V., see Castoldi, D. 413
Moretti, J.L., see Joulin, Y. 402
Morgan, H.W., see Patchett, M.L. 269
Mori, A., see Watanabe, Y. 373
Nagy, E.
—, Csipo, I., Degrell, I. and Szabo, G.
High-performance liquid chromatographic assay of 5-aminosalicylic acid and its acetylated metabolite in biological fluids using electrochemical detection 214
Nakanishi, Y., see Kojima, K. 203
Nambara, T., see Goto, J. 59
Ng, P., see Pereira, C.M. 208
Nolet, S., see Lemaire, S. 77
Novák, J., see Koupil, P. 99
Ostrovská, V., see Svobodová, X. 391
Pasutto, F.M., see Mehvar, R. 135
Patchett, M.L.
—, Monk, C.R., Daniel, R.M. and Morgan, H.W.
Determination of agmatine, arginine, citrulline and ornithine by reversed-phase liquid chromatography using automated pre-column derivatization with *o*-phthalaldehyde 269
Patrick, J.E., see Alton, K.B. 363
Peal, J.A., see Bailey, E. 187
Pechová, A., see Svobodová, X. 391
Pereira, C.M.
—, Tam, Y.K., Collins-Nakai, R.L. and Ng, P.
Simplified determination of captopril in plasma by high-performance liquid chromatography 208
Perry, H.E., see Schmid, B.J. 107
Petruzzi, R.F., see Alton, K.B. 363
Pfadenhauer, E.H.
High-performance liquid chromatographic assay for urinary metabolites of a drug after extraction by antiserum to the drug: an example using *erythro*-9-(2-hydroxy-3-nonyl)hypoxanthine 407
Philbert, M., see Bailey, E. 187
Platt, D., see Rieck, W. 121
Polacek, J.
—, Gunnarsson, P.O. and Brandin, S.
Determination of tauromustine, a nitrosourea-based antitumour agent, in plasma by high-performance liquid chromatography 424
Poláčiková, O., see Svobodová, X. 391
Porter, R.S., see Slegowski, M.B. 227
Prasad, V.K., see Carlin, A.S. 162
Przuntek, H., see Wallasch, Th.M. 175
Rapley, C.H., see Ubbink, J.B. 182
Rieck, W.
— and Platt, D.
High-performance liquid chromatographic method for the determination of α-amanitin and phalloidin in human plasma using the column-switching technique and its application in suspected cases of poisoning by the green species of amanita mushroom (*Amanita phalloides*) 121
Rose, L.M.
—, Tillery, K.F., El Dareer, S.M. and Hill, D.L.
High-performance liquid chromatographic determination of doxorubicin and its metabolites in plasma and tissue 419
Sager, A.O., see Carlin, A.S. 162
Sano, Y., see Goto, J. 59
Schatowitz, B.
— and Gercken, G.
Simultaneous determination of C_2–C_{22} non-esterified fatty acids and other metaboli-

cally relevant carboxylic acids in biological material by gas chromatography of their benzyl esters 257
Schmid, B.J.
—, Perry, H.E. and Idle, J.R.
Determination of nifedipine and its three principal metabolites in plasma and urine by automated electron-capture capillary gas chromatography 107
Schmid, K., see Graf, P. 353
Schmerr, M.J.F.
—, Goodwin, K.R. and Van Der Maaten, M.J.
High-performance gel-permeation chromatography of the surface glycoprotein from bovine leukemia virus 379
Schnell, A.M., see Ubbink, J.B. 182
Simmons, J.E., see Carlin, A.S. 162
Singh, A.K.
— and Ashraf, M.
Analysis of amino acids in brain and plasma samples by sensitive gas chromatography–mass spectrometry 245
Skelly, J.P., see Carlin, A.S. 162
Slegowski, M.B.
—, Miller, C. and Porter, R.S.
Simplified high-performance liquid chromatographic determination of flosequinan and its metabolite in plasma, serum and urine 227
Strecker, G., see Capon, C. 35
Sugi, H., see Watanabe, Y. 373
Sullivan, W.R.
— and Fox, K.E.
Gas chromatographic determination of the high-ceiling diuretic ethacrynic acid and its thiol adducts in plasma of guinea pigs 396
Suzuki, K.
—, Johno, I. and Kitazawa, S.
High-performance liquid chromatographic determination of nitrazepam in plasma and its application to pharmacokinetic studies in the rat 435
Svoboda, M., see Svobodová, X. 391
Svobodová, X.
—, Kováčová, D., Ostrovská, V., Pechová, A., Poláčiková, O., Kusala, S. and Svoboda, M.
Determination of nitroglycerin in human plasma using bonded-phase capillary column gas chromatography with electron-capture detection 391
Szabo, G., see Nagy, E. 214
Szathmary, S.Cs.
— and Luhmann, I.
Sensitive and automated gas chromatographic method for the determination of miconazol in plasma samples 193
Taha, A.M., see Marzouki, Z.M.H. 11
Takahashi, T., see Kojima, K. 203
Takahasi, H., see Matsumoto, K. 323
Tam, Y.K., see Pereira, C.M. 208
Tawa, R.
—, Koshide, K., Hirose, S. and Fujimoto, T.
Pre-column derivatization of sisomicin with *o*-phthalaldehyde-*β*-mercaptopropionic acid and its application to sensitive high-performance liquid chromatographic determination with fluorimetric detection 143
Tillery, K.F., see Rose, L.M. 419
Tofanetti, O., see Castoldi, D. 413
Tsuchiya, K., see Goto, J. 59
Ubbink, J.B.
—, Schnell, A.M. and Rapley, C.H.
Quantification of urinary xanthurenic acid excretion by anion-exchange solid-phase extraction and high-performance liquid chromatography 182
Uezono, Y., see Kojima, K. 203
Umino, M., see Matsumoto, K. 323
Van Der Maaten, M.J., see Schmerr, M.J.F. 379
Van der Walt, L.A.
— and Wittliff, J.L.
High-resolution separation of molybdate-stabilized progestin receptors using high-performance liquid chromatography 277
Vrbanac, J.J.
—, Eller, T.D. and Knapp, D.R.
Quantitative analysis of 6-keto-prostaglandin $F_{1\alpha}$ using immunoaffinity purification and gas chromatography–mass spectrometry 1
Wallasch, Th.M.
—, Henning, K., Lange, U., Kuhn, W., Eckhardt-Wallasch, H. and Przuntek, H.
Comparison of two methods for the determination of substance P immunoreactivity in human cerebrospinal fluid: sulphopropyl-Sephadex ion-exchange chromatography and/or reversed-phase high-performance liquid chromatography combined with radioimmunoassay 175
Walt, L.A. van der, see Van der Walt, L.A. 277
Watanabe, S., see Watanabe, Y. 373
Watanabe, Y.
—, Sugi, H., Watanabe, S. and Mori, A.
Identification of 1-guanidino-2-(4-imidazole)propionic acid in human urine by high-

performance liquid chromatography and field desorption mass spectrometry 373
Waud, W.R.
— and Hill, D.L.
High-performance liquid chromatographic determination of terephthalamidine in plasma 220
Weisweiler, P.
Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin–Sepharose chromatography 169
Wittliff, J.L., see Jung, S.S. 293
—, see Van der Walt, L.A. 277
Yee, G.C., see Gmur, D.J. 343
Zare, R.N., see Huang, X. 385
Zoest, A.R., see Hung, C.T. 331

Subject Index

Acetyl-5-aminosalicylic acid
HPLC assay of 5-aminosalicylic acid and its acetylated metabolite in biological fluids using electrochemical detection 214

cis-Aconitate
HPLC measurement of selected blood citric acid cycle intermediates 47

Adriamycin
HPLC determination of doxorubicin and its metabolites in plasma and tissue 419

Agmatine
Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 269

α-Amanitin
HPLC method for the determination of α-amanitin and phalloidin in human plasma using the column-switching technique and its application in suspected cases of poisoning by the green species of amanita mushroom (*Amanita phalloides*) 121

Amino acids
Analysis of amino acids in brain and plasma samples by sensitive GC–MS 245

Amino acids
Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 269

5-Aminosalicylic acid
HPLC assay of 5-aminosalicylic acid and its acetylated metabolite in biological fluids using electrochemical detection 214

Angiotensin converting enzyme inhibitor
Determination of the angiotensin converting enzyme inhibitor benazeprilat in plasma and urine by an enzymic method 353

Apolipoprotein E
Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin–Sepharose chromatography 169

Arginine
Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 269

Aromatic acids
Simultaneous determination of C_2–C_{22} non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 257

Benazeprilat
Determination of the angiotensin converting enzyme inhibitor benazeprilat in plasma and urine by an enzymic method 353

Benzoic acid
Determination of benzoic acid and hippuric acid in human plasma and urine by HPLC 67

Bile acid disulphates
Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 59

Bovine leukemia virus
High-performance gel-permeation chromatography of the surface glycoprotein from bovine leukemia virus 379

BTS 53554
Simplified HPLC determination of flosequinan and its metabolite in plasma, serum and urine 227

Busulfan
Determination of busulfan in human plasma by GC with electron-capture detection 303

Captopril
Simplified determination of captopril in plasma by HPLC 208

Carboxylic acids
Simultaneous determination of C_2–C_{22} non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 257

Chenodeoxycholate
Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 59

Cholate
Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 59

Cholesterol
Determination of coenzyme Q_{10}, α-tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 87

Citrate
HPLC measurement of selected blood citric acid cycle intermediates 47

Citric acid cycle intermediates
HPLC measurement of selected blood citric acid cycle intermediates 47
Citrulline
Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 269
Cloxacillin
Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 331
Coenzyme Q_{10}
Determination of coenzyme Q_{10}, α-tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 87
Corticoids
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 162
Corticosteroids
Determination of cortisol in human plasma by capillary GC–MS using [$^2H_5$]cortisol as an internal standard 237
Cortisol
Determination of cortisol in human plasma by capillary GC–MS using [$^2H_5$]cortisol as an internal standard 237
Cortisone
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 162
Cyclandelate
HPLC method for the determination of a cyclandelate metabolite, mandelic acid, in human plasma 203
Cyclosporine
Optimized HPLC method for the analysis of cyclosporine and three of its metabolites in blood and urine 233
Cyclosporins
HPLC column-switching method for two cyclosporine metabolites in blood 343
Debrisoquine
Comparison of GC and HPLC assays for the determination of debrisoquine and its 4-hydroxy metabolite in human fluids 311
Deoxycholate
Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 59
Dicarboxylic acids
Simultaneous determination of C_2–C_{22} non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 257
Dicloxacillin
Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 331
13-Dihydrodoxorubicin
HPLC determination of doxorubicin and its metabolites in plasma and tissue 419
Dilevalol
HPLC assay for dilevalol in human plasma and urine using a PRP-1 column and fluorimetric detection 363
1,3-Dinitrobenzene
Determination of 1,3-dinitrobenzene and its metabolites in rat blood by capillary GC with electron-capture detection 187
Doxorubicin
HPLC determination of doxorubicin and its metabolites in plasma and tissue 419
Ethacrynic acid
GC determination of the high-ceiling diuretic ethacrynic acid and its thiol adducts in plasma of guinea pigs 396
Fatty acid methyl esters
Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 11
Fatty acids
Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 11
Fatty acids
Thermospray–MS analysis of underivatized monohydroxy fatty acids: application to stimulated platelets 25
Fatty acids
Simultaneous determination of C_2–C_{22} non-esterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 257
Flosequinan
Simplified HPLC determination of flosequinan and its metabolite in plasma, serum and urine 227
Flucloxacillin
Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 331
Glycoproteins
Isolation of the major O-glycosidically linked oligosaccharides obtained by alkaline borohydride degradation of human meconium glycoproteins 35
Glycoproteins
High-performance gel-permeation chroma-

tography of the surface glycoprotein from bovine leukemia virus 379

1-Guanidino-2-(4-imidazole)propionic acid
Identification of 1-guanidino-2-(4-imidazole)propionic acid in human urine by HPLC and field desorption MS 373

Halothane
Comparison of the absolute calibration method with the method of standard addition for the determination of halothane in blood by GC headspace analysis 99

Hippuric acid
Determination of benzoic acid and hippuric acid in human plasma and urine by HPLC 67

Hydrocortisone
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 162

Hydroxy acids
Simultaneous determination of C_2–C_{22} nonesterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 257

4-Hydroxydebrisoquine
Comparison of GC and HPLC assays for the determination of debrisoquine and its 4-hydroxy metabolite in human fluids 311

Hydroxynimesulide
Simultaneous determination of nimesulide and hydroxynimesulide in human plasma and urine by HPLC 413

erythro-9-(2-Hydroxy-3-nonyl)hypoxanthine
HPLC assay for urinary metabolites of a drug after extraction by antiserum to the drug: an example using *erythro*-9-(2-hydroxy-3-nonyl)hypoxanthine 407

Isocitrate
HPLC measurement of selected blood citric acid cycle intermediates 47

Isoniazid
Determination of isoniazid in plasma by LC 441

Isopropyl-*p*-iodoamphetamine
Highly sensitive method for determination of isopropyl-*p*-iodoamphetamine by GC with electron-capture detection 402

Isoxazolyl penicillins
Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 331

Keto acids
Simultaneous determination of C_2–C_{22} nonesterified fatty acids and other metabolically relevant carboxylic acids in biological material by GC of their benzyl esters 257

α-Ketoglutarate
HPLC measurement of selected blood citric acid cycle intermediates 47

6-Keto-prostaglandin $F_{1\alpha}$
Quantitative analysis of 6-keto-prostaglandin $F_{1\alpha}$ using immunoaffinity purification and GC–MS 1

Lipids
Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 11

Lipoproteins
Determination of coenzyme Q_{10}, α-tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 87

Lipoproteins
Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin–Sepharose chromatography 169

Lithium
Quantitation of Li^+ in serum by capillary zone electrophoresis with an on-column conductivity detector 385

Mandelic acid
HPLC method for the determination of a cyclandelate metabolite, madelic acid, in human plasma 203

Miconazol
Sensitive and automated GC method for the determination of miconazol in plasma samples 193

Mitoxantron
Mitoxantron determination using HPLC: improved sensitivity by loop-column injection for dual-dose pharmacokinetic studies 429

Monohydroxy fatty acids
Thermospray–MS analysis of underivatized monohydroxy fatty acids: application to stimulated platelets 25

Nifedipine
Determination of nifedipine and its three principal metabolites in plasma and urine by automated electron-capture capillary GC 107

Nimesulide
Simultaneous determination of nimesulide and hydroxynimesulide in human plasma and urine by HPLC 413

Nitrazepam
HPLC determination of nitrazepam in plasma and its application to pharmacokinetic studies in the rat 435

3-Nitroacetanilide
Determination of 1,3-dinitrobenzene and its metabolites in rat blood by capillary GC with electron-capture detection 187
3-Nitroaniline
Determination of 1,3-dinitrobenzene and its metabolites in rat blood by capillary GC with electron-capture detection 187
Nitroglycerin
Determination of nitroglycerin in human plasma using bonded-phase capillary column GC with electron-capture detection 391
Nitrosourea
Determination of tauromustine, a nitrosourea-based antitumour agent, in plasma by HPLC 424
Oestrogen receptors
Influence of column lenght and pore size on HPIEC of estrogen and progestin receptors 293
Ofloxacin
Quantification of the enantiomers of ofloxacin in biological fluids by HPLC 153
Oligosaccharides
Isolation of the major O-glycosidically linked oligosaccharides obtained by alkaline borohydride degradation of human meconium glycoproteins 35
Organic acids
HPLC measurement of selected blood citric acid cycle intermediates 47
Ornithine
Determination of agmatine, arginine, citrulline and ornithine by reversed-phase LC using automated pre-column derivatization with *o*-phthalaldehyde 269
Oxaloacetate
HPLC measurement of selected blood citric acid cycle intermediates 47
Penicillins
Optimization of HPLC analysis for isoxazolyl penicillins using factorial design 331
Peptides
Sensitive method of detection, quantitation and purification of peptides using pre-column derivatization with phenyl isothiocyanate 77
Phalloidin
HPLC method for the determination of α-amanitin and phalloidin in human plasma using the column-switching technique and its application in suspected cases of poisoning by the green species of amanita mushroom (*Amanita phalloides*) 121
Prednisolone
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 162
Prednisolone acetate
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 162
Prednisone
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 162
Pristinamycin
Determination of PI_A and PII_A, the two main components of pristinamycin, by HPLC in plasma 197
Progestin receptors
High-resolution separation of molybdate-stabilized progestin receptors using HPLC 277
Progestin receptors
Influence of column length and pore size on HPIEC of estrogen and progestin receptors 293
Prostaglandins
Quantitative analysis of 6-keto-prostaglandin $F_{1\alpha}$ using immunoaffinity purification and GC–MS 1
Pyruvate
HPLC measurement of selected blood citric acid cycle intermediates 47
Sisomicin
Pre-column derivatization of sisomicin with *o*-phthalaldehyde–β-mercaptopropionic acid and its application to sensitive HPLC determination with fluorimetric detection 143
Steroids
Studies on steroids, CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 59
Steroids
Analysis of prednisolone acetate and related corticoids in swine plasma by reversed-phase HPLC 162
Steroids
Determination of cortisol in human plasma by capillary GC–MS using [$^2H_5$]cortisol as an internal standard 237
Substance P
Comparison of two methods for the determination of substance P immunoreactivity in human cerebrospinal fluid: sulphopropyl-Sephadex ion-exchange chromato-

graphy and/or reversed-phase HPLC combined with radioimmunoassay 175
Tauromustine
Determination of tauromustine, a nitrosourea-based antitumour agent, in plasma by HPLC 424
Terephthalamidine
HPLC determination of terephthalamidine in plasma 220
Theophylline
Automated determination of drugs in serum by column-switching HPLC. II. Separation of theophylline and its metabolites 323
Tiaprofenic acid
Rapid and sensitive HPLC assay of tiaprofenic acid enantiomers in human plasma and urine 135
α-Tocopherol
Determination of coenzyme Q_{10}, α-tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 87
Triglycerides
Fatty acid profiles of sebaceous triglycerides by capillary GC with mass-selective detection 11
Ursodeoxycholate
Studies on steroids. CCXXXIII. Separation and characterization of bile acid disulphates in human urine by HPLC 59
Very-low-density lipoproteins
Rapid quantification of apolipoprotein E enriched very-low-density lipoproteins by heparin–Sepharose chromatography 169
Vitamin E
Determination of coenzyme Q_{10}, α-tocopherol and cholesterol in biological samples by coupled-column LC with coulometric and UV detection 87
Xanthurenic acid
Quantification of urinary xanthurenic acid excretion by anion-exchange solid-phase extraction and HPLC 182